



## International Journal of Innovative Approaches in Agricultural Research

Volume 4, Issue 1 March 2020

[ijaar.penpublishing.net](http://ijaar.penpublishing.net)

ISSN: 2602-4772 (Online)

### Evaluation of Sweet Sorghum Biomass as an Alternative Livestock Feed

Celal Yücel

#### To cite this article

Yucel, C. (2020). Evaluation of Sweet Sorghum Biomass as an Alternative Livestock Feed. International Journal of Innovative Approaches in Agricultural Research, 4(1), 10-20. doi: 10.29329/ijaar.2020.238.2

Published Online	March 29, 2020
Article Views	67 single - 87 cumulative
Article Download	280 single - 553 cumulative
DOI	<a href="https://doi.org/10.29329/ijaar.2020.238.2">https://doi.org/10.29329/ijaar.2020.238.2</a>

Pen Academic is an independent international publisher committed to publishing academic books, journals, encyclopedias, handbooks of research of the highest quality in the fields of Education, Social Sciences, Science and Agriculture. Pen Academic created an open access system to spread the scientific knowledge freely. For more information about PEN, please contact: [info@penpublishing.net](mailto:info@penpublishing.net)





Original article

## Evaluation of Sweet Sorghum Biomass as an Alternative Livestock Feed

Celal Yücel \*

Department of Field Crops, Faculty of Agricultural, Sirnak University, Sirnak, Turkey

### Abstract

In case of using sorghum biomass as a source of feed, quality attributes of different forms of the biomass should be known. This study was conducted to determine quality attributes of sweet sorghum (*Sorghum bicolor var. saccharatum (L.) Mohlenbr.*) biomass in different forms. Six different sweet sorghum varieties (M81-E, Ramada, Roma, Topper 76, UNL Hybrid and No 91) were used in different forms. Quality attributes of sweet sorghum herbage before silage (SSH), whole plant silage (WPS) and sap-extracted bagasse silage (SES) were determined. Harvests were performed at milk-dough stage of the plants. While green herbage had greater crude protein ratios, whole plant silages had greater values for the other quality attributes. The sap-extracted bagasse silage had a mean RFV of over 111 indicating a high quality feed. As compared to other varieties, Topper 76 and Ramada varieties had greater values for quality attributes of sweet sorghums.

**Keywords:** Sweet Sorghum, Silage, Biomass, Quality, Feed.

**Received:** 07 October 2019 \* **Accepted:** 23 November 2019 \* **DOI:** <https://doi.org/10.29329/ijjaar.2020.238.2>

### \* Corresponding author:

Celal Yücel is an professor in the Department of Field Crops at Sirnak University in Turkey. Her research interests include the forage crops and energy crops. He has lived, worked, and studied in Sirnak, Turkey.  
Email: celalyucel1@gmail.com

## **INTRODUCTION**

As compared to the other feed crops, sweet sorghum has less fertilizer demands and soil preference. Sweet sorghum is quite resistant to draughts and high temperatures (Mastrorilli et al., 1999; Gnansounou et al., 2005; Tesso et al., 2005), more tolerant to floods and salinity (Almodares et al., 2008). Sorghum plants have also quite high energy contents (Negro et al., 1999). Although the water demand of sorghum is lower than those of maize and sugar cane, it has greater potential of biomass than many other C4 plants (Gnansounou et al., 2005). Sweet sorghum is a short-day plant and majority of sorghum varieties require very high temperatures and achieve the best growth and development under these conditions (Reise and Almodares, 2008). Sweet sorghum stalks are ensiled and thus generate a quality feed source for ruminants (Jafarina et al., 2005). According to Kumari et al. (2013), sweet sorghum bagasse, which is an agricultural byproduct, may constitute a good source of silage without any additives and reliably be used in animal diets. Naeini et al. (2014) reported that addition of substances such as urea and molasses into sorghum silage may improve the quality of the silage even further. Many new sorghum varieties have been developed for silage purposes in recent years. They provide greater biomass yield per unit area and they have almost equivalent quality attributes to maize silage (Yucel et al., 2018). Production of summer C4 plants like sorghum with high yield potential per unit-area should be increased to provide quality roughage source for livestock. Number of studies about the potential use of sweet sorghum as a feed source for ruminants in Turkey is quite limited. Thus, information about potential uses of sweet sorghum plants is not available.

This study was conducted to determine the forage quality of six sorghum genotypes grown under the ecological conditions of the Mediterranean Region of Turkey. Whole plants (leaf-stem-panicle) and sap-extracted plants (bagasse) were ensiled and silage quality attributes were also determined in this study.

### **Materials and Methods**

#### ***Materials***

The material included M81-E, Ramada, Roma and Topper 76 varieties, and UNL-hybrid (26297 X M81 E) of sweet sorghum obtained from the University of Nebraska (USA) in addition to the populations of no91 with USDA-Taiwan origins.

#### ***Method***

Field experiments were conducted over the experimental fields of Eastern Mediterranean Agricultural Research Institute (Dogankent-Adana/TURKEY) (36° 51' 35" N and 35° 20' 43" E) in 2016 under second crop conditions in split-plots experimental design with 4 replications. Sowing was performed on 13<sup>th</sup> of June, 2016 following the harvest of wheat. Before sowing, 50 kg/ha of nitrogen and phosphorus were applied as base fertilizer. All varieties were manually sown at 70 cm row spacing

and 15 cm on-row plant spacing. Each plot had 4 rows of 5 m long. When the plants reached to 40-50 cm heights, 50 kg/ha of pure nitrogen was applied as top-dressing fertilizer. Harvesting was carried out between the milk and dough stage of the plants. A total of 15 plants were randomly selected from each plot: 5 plants for herbage, 5 plants for whole plant silage and 5 plants for sap-extracted plant silage.

Three different treatments were conducted to determine the quality attributes of sweet sorghum varieties:

1. Quality values of the herbage material (pre-ensilage): 5 plants harvested from the field were cut into pieces of 15-20 cm, they were put into paper bags, dried in an oven at 65 °C until a constant weight for about 7-10 days. They were ground to pass 1-2 mm sieves.
2. Quality values of whole plant (leaf-stem-panicle) silage: 5 plants harvested from the field were chopped into 3-5 cm pieces in a chipper-chopper machine. Chopped samples were then used to make whole plant silage.
3. Quality values of sap-extracted (squeezed-bagasse) silage: Stalks of 5 plants were squeezed by a specially designed device to extract the plant sap, then resultant bagasse material was chopped into 3-5 cm pieces with a grinding machine. Chopped samples were used to make sap-extracted plant silage (Yücel et al., 2017).

About 1-kg green samples were taken from chopped silage materials, green samples were put into special plastic bags (110 microns of thickness or thicker), 99.9% of the oxygen inside the bags was removed by a Crompack vacuum device, and the bags were sealed automatically to finalize the silage process. Ensiled material was labeled and kept at room temperature (17-25 °C) for 60 days. For all treatments, silage samples were dried, weighed and ground to pass 1-2 mm sieves. The Kjeldahl method was used to determine the nitrogen (N) contents of the samples. The crude protein ratio was determined by the formula  $N \times 6.25$  (AOAC 1990). The % NDF and ADF contents which constituted the cell membrane components of the feeds were determined by the methods described by Van Soest et al. (1991) in an ANKOM fiber analysis device (Fiber Analyzer). Digestible dry matter (DDM) ratio, dry matter intake (DMI) and relative feed value (RFV) were calculated by the formulae reported by Schroeder (1994). Accordingly,  $DDM = 88.9 - (0.779 \times \% ADF)$ ;  $DMI = 120 / \% NDF$ ;  $RFV = (\% DDM \times \% DMI) / 1.29$ . Net energy-lactation (NEL-Mcal/kg) was calculated by the formula:  $NE = 1.892 - (0.0141 \times ADF)$  (Anonymous, 2018).

JUMP statistical software was used for statistical analyses. Significant means were compared with the aid of LSD test at  $P < 0.05$  level.



**Figure 1. a)**

**b)**

**c)**

a) SSH: Sweet sorghum herbage b) SES: Sap-extracted plant silage c) WPS: Whole plant silage

## **Results and Discussion**

### ***Crude Protein Ratio (%)***

There were significant differences in crude protein ratios of both the varieties and the treatments (Table 1). The crude protein ratios of the treatments and varieties varied between 2.53 - 5.99%. The greatest crude protein ratio of the varieties (5.08%) was obtained from Ramada variety (Table 1). The greatest crude protein ratio of the treatments (4.88%) was obtained from sweet sorghum herbage and it was respectively followed by whole plant silage (4.06%) and sap-extracted plant (bagasse) silage (3.65%) treatments. The quality values of green herbage and whole plant silage were higher than those of sap-extracted plant silage since whole plants contained stalks, leaves and panicles, and panicle contained grains. As the grains in the panicles of the ensiled plants were in dough stage, they provided quite high protein values. CP contents of sorghum grains were reported as between 7 - 15% (FAO 1995; Beta et al., 1995). CP ratios of sorghum herbage were reported as between 4.10-8.01% (Rodrigues et al., 2006; Junior et al., 2015; Durul, 2016), and the crude protein ratios of silage sorghum were reported as between 5.6-11.71% (Mahmood et al., 2013 a,b; Karadağ and Özkurt, 2014; Neto et al., 2017). It was reported in previous studies that CP ratios of sap-extracted sweet sorghum silage varied between 4.1 - 7.26% (Kumari et al., 2013; Trulea et al., 2013; Naeini et al., 2014; Rocha et al., 2018) and the CP ratio of silage made out of the bagasse of sweet sorghum that was squeezed with its leaf was 7.48% (Vidya et al., 2016).

### ***Crude Ash (CA, %)***

There were significant differences in crude ash contents of the varieties and the treatments. Crude ash contents of the treatments and varieties varied between 4.20 - 7.38%. The greatest crude ash content was obtained from whole plant silage and it was followed by the other treatments (Table 1). Crude ash contents of sweet sorghum varieties were reported as between 6.15-13.08% (Chakravarthi et al., 2017); they had a crude ash quantity of 79.3 g in 1 kg of DM (Cattani et al. 2017); CA content of herbage before

silage was 3.35% and CA content of sorghum silage was 3.54%, while the CA content of sorghum bagasse before silage was 2.52%, the CA content of bagasse silage was 2.55% (Trulea et al. 2013); the crude ash contents of the green material of sorghum and the bagasse of sorghum varied as between 59 - 60 g per 1 kg of DM, and the crude ash values of the ensiled sorghum and bagasse were found respectively as 57 and 70 g per 1 kg of DM (Naeini et al., 2014).

**Table 1.** Mean Crude Protein and Crude Ash Ratios of Sweet Sorghum Biomass Based on Different Treatment

Cultivars	Crude Protein Ratio (%)				Ash Content (%)			
	SSH*	WPS	SES	Mean	SSH*	WPS	SES	Mean
M81-E	4.56	3.36	2.53	<b>3.50 C<sup>+</sup></b>	5.42	6.56	6.18	<b>6.05</b>
Ramada	5.99	5.01	4.25	<b>5.08 A</b>	6.27	6.25	6.28	<b>6.27</b>
Roma	5.13	4.46	4.29	<b>4.63 B</b>	6.13	7.38	6.57	<b>6.69</b>
Topper 76	4.14	3.77	3.47	<b>3.79 C</b>	5.64	7.21	4.20	<b>5.66</b>
UNL Hybrid	5.04	4.21	4.04	<b>4.43 B</b>	6.35	6.60	5.05	<b>6.00</b>
No91	4.42	3.55	3.24	<b>3.74 C</b>	5.20	5.67	5.14	<b>5.34</b>
<b>Mean</b>	<b>4.88 A</b>	<b>4.06 B</b>	<b>3.65 C<sup>1</sup></b>		<b>5.84 B</b>	<b>6.60 A</b>	<b>5.57 B<sup>1</sup></b>	
CV (%)	9.48				19.48			

\*) SSH: Sweet sorghum herbage      b) WPS: Whole plant silage      c) SES: Sap-extracted plant silage

+) There is no statistically significant difference between the mean values that are represented by the same letter on the same column according to the LSD test at P≤0.05

1) There is no statistically significant difference between the mean values that are represented by the same letter on the same row according to the LSD test at P≤0.05

### **Acid Detergent Fiber (ADF, %)**

There were significant differences in acid detergent fiber ratios of the treatments, varieties and treatment x variety interactions (Table 2). The ADF values of different variety-treatment combinations varied between 25.00-39.57%. The silage made out of the whole plant had lower ADF values. It may be due to the silages made out of the whole plant contained both the leaves and panicles. Moreover, leaves were found to contain lower levels of cell-membrane materials as compared to stalks. It was reported that cell-membrane components such as cellulose and lignin were found at higher rates in the stalks as compared to the leaves (Aman, 1993) and these components negatively correlated with digestion (Hatfield, 1993). Among the varieties, Topper 76 had the lowest ADF content. The ADF content of sweet sorghum silages were reported as between 27.9-42.0% (Mahmood et al., 2013; Durul, 2016). It was determined that the ADF content of silage made out of the bagasse of squeezed sweet sorghum varied between 19.16-46.82% (Ávila et al., 2013; Kumari et al., 2013; Rocha et al., 2018), and the ADF content of silage made out of the bagasse of sweet sorghum that is squeezed with its leaves was 46.75% (Vidya et al., 2016). Naeini et al. (2014) reported the ADF values of the green materials of maize, sorghum and sorghum bagasse respectively as 263, 213 and 258 g per 1 kg of DM and ADF values of silage made out of these materials respectively as 272, 254 and 246 g per 1 kg of DM. It was reported

that sorghum and sorghum bagasse without any additives had lower NDF and ADF values than maize (Naeini et al., 2014).

### **Neutral Detergent Fiber (NDF, %)**

Neutral detergent fiber ratios of the treatments, varieties and treatment x variety interactions are provided in Table 2. The NDF values of different treatment-variety combinations varied between 42.58-60.70%. The silage made out of bagasse had lower NDF values. Since the silages made out of bagasse did not contain leaves and panicles, some quality parameters may be expected to be low. The Topper 76 variety had the lowest NDF value. The mean NDF values of sweet sorghum varieties were reported as between 32.6-54.9 % (Gomes et al., 2006; Machado et al., 2012; Durul, 2016; Neto et al., 2017). Karadağ and Özkurt (2014) reported NDF ratios of silage sorghum varieties as between 61.23-63.00%. The NDF ratios of silage made out of the bagasse of squeezed sweet sorghum was reported as between 39.23 - 75.4% (Ávila et al., 2013; Kumari et al., 2013; Rocha et al., 2018), and the NDF content of silage made out of the bagasse of sweet sorghum that was squeezed with its leaves was 71.81% (Vidya et al., 2016). Naeini et al. (2014) reported the NDF values of the green materials of maize, sorghum and sorghum bagasse respectively as 526, 447 and 491 g per 1 kg of DM.

**Table 2.** The Mean ADF and NDF Values in Different Varieties of Sorghum and Different Treatment

Cultivars	Acid Detergent Fiber (%)				Neutral Detergent Fiber (%)			
	SSH*	WPS	SES	Mean	SSH*	WPS	SES	Mean
M81-E	30.35 cde <sup>2</sup>	29.18 c-f	39.57 a	<b>33.05 A<sup>+</sup></b>	47.64 c-g	46.89 d-h	57.59 a <sup>2</sup>	<b>50.71 A<sup>+</sup></b>
Ramada	29.29 cde	25.00 g	30.35 cde	<b>28.45CD</b>	48.10 b-f	43.29 gh	48.56 b-f	<b>46.65BC</b>
Roma	27.88 d-g	26.25 fg	35.48 b	<b>29.87 BC</b>	45.63 e-h	42.83 h	52.40 b	<b>46.95 BC</b>
Topper 76	25.11 g	26.19 fg	30.73 cd	<b>27.34 D</b>	43.39 gh	46.22 d-h	44.02 fgh	<b>44.54 C</b>
UNL H.	32.83 bc	25.76 fg	36.05 ab	<b>31.55 AB</b>	50.79 bcd	42.58 h	52.13 bc	<b>48.50 AB</b>
No91	30.81 cd	26.69 efg	39.50 a	<b>32.34 A</b>	48.86 b-e	44.05 fgh	60.70 a	<b>51.20 A</b>
<b>Mean</b>	<b>29.50 B</b>	<b>26.51 C</b>	<b>35.28 A<sup>1</sup></b>		<b>47.40 B</b>	<b>44.31 C</b>	<b>52.56 A<sup>1</sup></b>	
CV (%)	8.54				6.86			

\*) **SSH:** Sweet sorghum herbage, **WPS:** Whole plant silage, **SES:** Sap-extracted plant silage

+) There is no statistically significant difference between the mean values that are represented by the same letter on the same column according to the LSD test at  $P \leq 0.05$

1) There is no statistically significant difference between the mean values that are represented by the same letter on the same row according to the LSD test at  $P \leq 0.05$

2) There is no statistically significant difference between different cultivars-treatment combination that are represented by the same letter according to the LSD test at  $P \leq 0.05$ .

### Dry Matter Intake (DMI, %)

There were significant differences in dry matter intake ratios of the treatments, varieties and treatment x variety interactions (Table 3). The DMI ratios of different variety-treatment combinations varied between 2.39-2.70% . The DMI ratio of silage made out of the whole plant was significantly greater than those of the other treatments. The Topper 76 variety had the greatest dry matter intake ratio. Lower DMI ratios of sap-extracted bagasse silages were attributed to high NDF values of these silages made out of only the stalks without the leaves and panicles (grains). It is known that there was a negative correlation between NDF and DMI. Karthikeyan et al. (2017) reported dry matter intake ratios of sorghum varieties as between 1.67-2.20 % with an average value of 1.93 %.

**Table 3.** The Mean DMI Ratio and DDMR Values in Different Varieties of Sorghum and Different Treatment

Cultivars	Dry Matter Intake (%)				Digestible Dry Matter Ratio (%)			
	SSH*	WPS	SES	Mean	SSH*	WPS	SES	Mean
M81-E	2.52 c-f <sup>2</sup>	2.56 b-e	2.09 g-h	<b>2.39 C<sup>+</sup></b>	65.23 cde	66.17 b-e	58.08 g <sup>2</sup>	<b>63.16 D<sup>+</sup></b>
Ramada	2.50 def	2.80 a	2.48 def	<b>2.59 AB</b>	65.54 cde	69.43 a	62.26 cde	<b>66.74 AB</b>
Roma	2.64 a-d	2.81 a	2.30 fg	<b>2.58 AB</b>	67.18 a-d	68.45 ab	61.26 f	<b>65.63 BC</b>
Topper 76	2.77 ab	2.61 a-e	2.74 abc	<b>2.70 A</b>	69.34 a	68.50 ab	64.96 de	<b>67.60 A</b>
UNL H.	2.39 ef	2.83 a	2.31 fg	<b>2.51 BC</b>	63.33 ef	68.83 ab	60.82 fg	<b>64.32 CD</b>
No91	2.46 def	2.73 abc	1.99 h	<b>2.39 C</b>	64.90 de	68.10 abc	58.13 g	<b>63.71 D</b>
<b>Mean</b>	<b>2.55 B</b>	<b>2.72 A</b>	<b>2.32 C<sup>1</sup></b>		<b>65.92 B</b>	<b>68.25 A</b>	<b>61.42 C<sup>1</sup></b>	
CV (%)	6.40				3.11			

\*) SSH: Sweet sorghum herbage, WPS: Whole plant silage, SES: Sap-extracted plant silage

+) There is no statistically significant difference between the mean values that are represented by the same letter on the same column according to the LSD test at P≤0.05

1) There is no statistically significant difference between the mean values that are represented by the same letter on the same row according to the LSD test at P≤0.05

2) There is no statistically significant difference between different cultivars-treatment combination that are represented by the same letter according to the LSD test at P≤0.05.

### Digestible Dry Matter (DDM, %)

There were significant differences in digestible dry matter ratios of the treatments, varieties and treatment x variety interactions (Table 3). The DDM ratios of different variety-treatment combinations varied between 58.08-69.43%. The DDM ratio of silage made out of the whole plant was greater than the other treatments. The Topper 76 variety had the lowest DDM ratio. Increasing sugar content of sorghum stalks increases digestibility and feed quality (Poehlman, 1994; Blümmel et al., 2009). The digestible dry matter ratios of sorghum silages were reported as between 57.02-66.30% (Junior et al., 2015; Karthikeyan et al., 2017).

### Relative Feed Value

There were significant differences in relative feed values of the treatments, varieties and treatment x variety interactions (Table 4). The RFV of the varieties varied between 81.5- 146.9. The greatest value was obtained from Topper 76 variety and the lowest value was obtained from M81-Evariety. The values of the other genotypes were in between them. The RFV of alfalfa is assumed to be 100 when the plants were harvested at 100% flowering stage. Durul (2016) reported RFV of sweet sorghum silages as between 104-126.

**Table 4.** The Mean RFV and NEL Values in Different Varieties of Sorghum and Different Treatment

Cultivars	Relative Feed Value				Net Energy Lactation (%)			
	SSH*	WPS	SES	Mean	SSH*	WPS	SES	Mean
M81-E	127.4 cd <sup>2</sup>	131.5 bcd	94.4 f	<b>117.8 C<sup>+</sup></b>	1.849 a	1.852a	1.334 d	<b>1.678B<sup>+</sup></b>
Ramada	127.3 cd	150.7 a	125.5 cd	<b>134.5 AB</b>	1.850 a	1.856 a	1.464 b	<b>1.723 A</b>
Roma	137.3 abc	148.9 a	109.6 ef	<b>131.9 B</b>	1.853 a	1.855 a	1.392 c	<b>1.700AB</b>
Topper 76	148.8 a	138.5 abc	137.9 abc	<b>141.7 A</b>	1.857 a	1.854 a	1.459 b	<b>1.723 A</b>
UNL Hybrid	117.8 de	151.4 a	108.8 ef	<b>126.0 BC</b>	1.846 a	1.854 a	1.384 c	<b>1.695 B</b>
No91	123.7 cde	144.4 ab	90.2 g	<b>119.4 C</b>	1.849 a	1.855 a	1.335 d	<b>1.679 B</b>
<b>Mean</b>	<b>130.4 B</b>	<b>144.2 A</b>	<b>111.1 C<sup>1</sup></b>		<b>1.850 A</b>	<b>1.854 A</b>	<b>1.395 B<sup>1</sup></b>	
CV (%)	8.63				1.79			

\*) **SSH:** Sweet sorghum herbage, **WPS:** Whole plant silage, **SES:** Sap-extracted plant silage

+ ) There is no statistically significant difference between the mean values that are represented by the same letter on the same column according to the LSD test at P≤0.05

1) There is no statistically significant difference between the mean values that are represented by the same letter on the same row according to the LSD test at P≤0.05

2) There is no statistically significant difference between different cultivars-treatment combination that are represented by the same letter according to the LSD test at P≤0.05

### Net Energy - Lactation (Mcal/kg)

There were significant differences in net energy values of the treatments, varieties and treatment x variety interactions (Table 4). Net energy values of the varieties varied between 1.273 - 1.473 Mcal kg<sup>-1</sup> with the greatest value from Topper 76 variety and lowest value from no91 variety. It was stated that the energy content of sweet sorghum was high due to its high water-soluble carbohydrate contents (Kaiser et al., 2004). Cattani et al. (2017) reported NE contents of sorghum silages as between 1.82-0.92 Mcal per kg of DM.

### Conclusion

Quality attributes of sorghum biomass in different forms (as fresh herbage, whole plant silage, sap-extracted bagasse silage) were analyzed in this study. While green herbage had greater crude protein ratios, whole plant silages had greater values for the other quality attributes. RFV was calculated by

taking 100% flowering stage of alfalfa into consideration. Accordingly, the sap-extracted bagasse silage had a mean RFV of over 111, the values of Topper 76 and Ramada varieties were even higher, respectively as 137.9 and 125.5, and these values were also higher than those of several other feed crops. In terms of the varieties, Topper 76 and Ramada showed higher values of quality.

## REFERENCES

- Almodares, A., M.R. Hadi and H. Ahmadpour (2008). Sorghum stem yield and soluble carbohydrates under phenological stages and salinity levels. *Afr. J. Biotechnol.*, **7**, 4051-4055.
- Aman, P. (1993). Composition and structure of cell wall polysaccharides in forages. In: Jung HG, Buxton DR, Hatfield RD, Ralph J. (Eds.), *Forage Cell Wall Structure and Digestibility*. ASA, CSSA, and SSSA, Madison, WI, pp. 183-199.
- Anonymous, (2018). SGS Agrifood Laboratories. Feed/Forages Calculations. [http://www.agtest.com/articles/feed%20and%20forages%20calculations\\_new.pdf](http://www.agtest.com/articles/feed%20and%20forages%20calculations_new.pdf). AOAC, Association of Official Analytical Chemists (1990). *Official method of analysis. 15th ed.* Washington, DC. USA, 66-88.
- Avila, S.C., A.A. Martins, G. V. Kozloski, T. Orlandi, M. P. Mezzoma, C.M. Stefanello, F. Hentz, and P. Castagnino (2013). Sunflower meal supplementation to wethers fed sorghum bagasse silage. *Ciênc. Rural*, **42** (7), 1245-1250.
- Beta, T., L.W. Rooney and R.D. Waniska (1995). Malting characteristics of sorghum cultivars. *Cereal Chem.*, **72**, 533-538.
- Blümmel, M., S.S. Rao, S. Palaniswami, L. Shahand B.V.S. Reddy (2009). Evaluation of sweet sorghum (*Sorghum bicolor* (L.) Moench) used for bio-ethanol production in the context of optimizing whole plant utilization. *Anim. Nutr. Feed Techn.*, **9**, 1-10.
- Cattani, M., N. Guzzo, R. Mantovani and L. Bailoni (2017). Effects of total replacement of corn silage with sorghum silage on milk yield, composition, and quality. *J. Anim. Sci. Biotechnol.*, **8**, 15. DOI 10.1186/s40104-017-0146-8.
- Chakravarthi, M.K., Y.R. Reddy, K.S. Rao, A. Ravi, B. Punyakumari and B. Ekambaram (2017). A study on nutritive value and chemical composition of sorghum fodder. *Int. J. Environ. Sci. Te.*, **6** (1), 104-9.
- Durul, G. (2016). Farklı biçim zamanlarının tatlı sorgum (*sorghum bicolor* (L.) moench var. *saccharatum*) ve fasulye (*Phaseolus vulgaris*) silaj karışımlarında bazı kalite özelliklerine etkisi. *Ege Üni. Fen Bil. Ens., Tarla Bitk.ABD, Y. Lisans Tezi*, 72 s.
- FAO, Food and Agricultural Organization. (1995). Sorghum and millet in human nutrition. *FAO Food and Nutrition Series No. 27*. ISBN 92-5-103381-1. Consulted on 10 September 2005 at: <http://www.fao.org>
- Gnansounou, E., A. Dauriat and C.E. Wyman (2005). Refining sweet sorghum to ethanol and sugar: economic trade-offs in the context of North China. *Bioresour Technol.*, **96**, 985-1002.
- Gomes, S.O., J.B. Pitombeira, J.N.M. Neiva and M.J.D. Candidado (2006). Agronomic behavior and forage composition of sorghum cultivars in the State of Ceará. *Rev. Ciênc. Agron.*, **37** (2), 221-2277.

- Hatfield, R.D. (1993). Cell wall polysaccharide interactions and digestibility. In 'Forage cell wall structure and digestibility'. (Eds HG Jung, DR Buxton, RD Hatfield, J Ralph) pp. 285–313. (ASA, CSSA, and SSSA: Madison, WI).
- Jafarinia, M., A. Almodares and M. Khorvash (2005). Using sweet sorghum bagasse in silo In: Proceeding of the 2nd Congress of Using Renewable Sources and Agric. Wastes (Eds. M Jafarinia, A Almodares & M Khorvash). Khorasgan Azade University, Isfahan, Iran.
- Junior, M.A.P.O., M. Retore, D.M. Manarelli, F.B. de Souza, L.M. Ledesma and A.C.A. Orrico (2015). Forage potential and silage quality of four varieties of saccharine sorghum. *Pesqui. Agropecu. Bras.*, 50 (12), 1201-1207.
- Kaiser, A.G., J.W. Plitz, H.M. Burns and N.W. Griffiths (2004). Successful Silage. Dairy Australia NSW Department of Primary Industries, 468p.
- Karadağ, Y. and M. Özkurt (2014). İkinci ürün olarak yetiştirilebilecek silajlık sorgum (*Sorghum Bicolor* (L). Moench) çeşitlerinde farklı sıra aralıklarının verim ve kalite üzerine etkisi. *Gaziosmanpaşa Üni. Ziraat Fak. Derg.*, 31(1), 19-24.
- Karthikeyan, B. J., C. Babu and J. J. Amalraj (2017). Nutritive value and fodder potential of different sorghum (*Sorghum bicolor* L. Moench) cultivars. *Int. J. Curr. Microbiol. App. Sci.*, 6(8), 898-911.
- Kumari, N. N., Y.R. Reddy, M. Blümmel, D. Nagalakshmi and T. Monica (2013). Effect of feeding sweet sorghum bagasse silage with or without chopping on nutrient utilization in deccani sheep. *Anim. Nutr. Feed Techn.*, 13, 243-249.
- Machado, L.C, W.M. Ferreira and C. Scpinello (2012). Apparent digestibility of simplified and semi-simplified diets, with and without addition of enzymes, and nutritional value of fibrous sources for rabbits. *R. Bras. Zootec.*, 41, 1662-1670.
- Mahmood, A., H. Ullah, M. Ijaz, M.M. Javaid, A.N. Shahzad and B. Honermeier (2013a). Evaluation of sorghum hybrids for biomass and biogas production. *Aust. J. Crop Sci.*, 7 (10), 1456-1462.
- Mahmood, A., H. Ullah, A.N. Shahzad, H. Ali, S. Ahmad, M. Zia-Ul-haq, B. Honermeier and M. Hasanuzzaman (2013b). Dry Matter Yield and Chemical Composition of Sorghum Cultivars with Varying Planting Density and Sowing Date. *Sains Malays.*, 42(10), 1529–1538.
- Mastrorilli, M., N. Katerji and G. Ranai (1999). Productivity and water use efficiency of sweet sorghum as affected by soil water deficit occurring at different vegetative growth stages. *Eur. J. Agron.*, 11, 207-215.
- Naeini, S. Z., M. Khorvash, E. Rowghani, A. Bayat and Z. Nikousefat (2014). Effects of urea and molasses supplementation on chemical composition, protein fractionation and fermentation characteristics of sweet sorghum and bagasse silages as alternative silage crop compared with maize silage in the arid areas. *Res. Opin. Anim. Vet. Sci.*, 4(6), 343-352.
- Negro, M.J., M. L. Solano, P. Ciria and J. Carrasco (1999). Composting of sweet sorghum bagasse with other wastes. *Bioresour Technol.*, 67, 89-92.
- Neto, A.B., R. H. Pereira dos Reis, S. Cabral L. da, J. G. Abreu, D.P. Sousa and F.G. Sousa (2017). Nutritional value of sorghum silage of different purposes. *Ciênc. Agrotec.*, 41(3), 288-299.
- Poehlman, J. M (1994). Breeding sorghum and millet. In *Breeding field crops*, 3rd ed, ed. J.M. Poehlman, 508-541. Ames, Iowa, USA: Iowa State University Press.

- Reisi, F. and A. Almodares (2008). The effect of planting date on amylose content in sorghum and corn. In: Proceeding of the 3rd Int. Biol. conference (Eds. F Reisi & A Almodares). Tehran, Iran.
- Rocha, G., F. Mara and E.A. Ricardo et al. (2018). Fermentation characteristics and bromatological composition of sweet sorghum bagasse silages. Rev. Bras. de Saude e Prod. Anim., 19 (2), 157-165.
- Rodrigues, F.O., A. F. de S. França R. de P. Oliveira, E.R. de Oliveria, B. Rosa, T. V. Soares and S. Q.S Mello (2006). Produção e composição bromatológica de quatro híbridos de sorgo forrageiro (*Sorghum bicolor* (L.) Moench) submetidos através doses de nitrogênio. Ciênc. Anim. Bras., 7, 37-48.
- Schroeder, J. W. (1994). Interpreting forage Analysis. Extention Dairy specialist (NDSU). AS-1080, North Dakota State University.
- Tesso, T. T., L. E. Clafin and M.R. Tuinstra (2005). Analysis of stalk rot resistance and genetic diversity among drought tolerant sorghum genotypes. Crop Sci., 45, 645-652.
- Trulea, A., T. Vintila, G. Pop, R. Sumalan and S. Gaspar (2013). Ensiling sweet sorghum and maize stalks as feedstock for renewable energy production. Res. J. Agric. Sci., 45, 193-199.
- Van Soest, P.J., J. D. Robertson and B. A. Lewis (1991). Methods for dietary fibre, neutral detergent fibre and non-starch polysaccharides in relation to animal nutrition. J. Dairy Sci., 74, 3583-3597.
- Vidya B, Reddy Y R, Rao D S, Reddy V R, Kumari N N & Blummel M (2016). Effect of supplementation of concentrate to sweet sorghum bagasse with leaf residue silage on nutrient utilization and nitrogen balance in native sheep. *Indian J. Anim. Res.*, 50 (3), 387-391.
- Yücel, C., İ. İnal, F. Gündel, D. Yücel, A. Aktaş, H.A. Karaağaç, R. Hatipoğlu. and I. Dweikat (2017). Biyoetanol Üretiminde Kullanılmış Tatlı Sorgum Saplarının Silaj Kalite Özelliklerinin Belirlenmesi. KSÜ Doğa Bil. Derg., 20 (Özel Sayı), 144-148.
- Yücel, C., R. Hatipoğlu, I. Dweikat, I. İnal, F. Gündel and H. Yucel (2018). A Research on Determination of Biyo- etanol Production Potential of Different Sweet Sorghum (*Sorghum bicolor* var. *saccharatum* (L.) Mohlenbr.) Genotypes in Çukurova and GAP Regions. TUBITAK 1003 Project Final Report.